

Challenges with Anaesthesia During Liposuction

Jacob Taylor[†]

ABSTRACT

One of the most common treatment options in aesthetic surgery, liposuction has certain special anaesthetic considerations. Liposuction is frequently carried out at a doctor's office. Based on the amount of infiltration or wetting solution injected, there are four basic types of liposuction techniques: dry, wet, superwet, and tumescent procedure. One of the most popular liposuction methods is known as the tumescent technique, which involves injecting large amounts of diluted local anaesthetic (wetting solution) into the fat in order to facilitate anaesthesia and reduce blood loss. The injection of lignocaine may contain a significant amount between 35 mg/kg and 55 mg/kg raising questions about the toxicity of the local anaesthetic.

Keywords: Anaesthesia; Aesthetic surgery; Liposuction; Blood

Introduction

According to the amount of solution aspirated, there are two types of liposuction: high volume (>4,000 ml aspirated) and low volume (4,000 ml aspirated). Large-volume liposuction necessitates general anaesthesia, whereas small-volume liposuction can be performed under local/monitored anaesthesia care. The intraoperative fluid management, together with hemodynamic monitoring and temperature control, must all be carefully titrated as a significant volume of wetting solution is injected into the subcutaneous tissue. Since the blood is mingled with the inhaled fat, measuring the blood loss is challenging. The majority of obese patients choose liposuction as a rapid weight loss procedure, thus all obesity-related issues must be addressed in the preoperative evaluation.

is Liposuction common plastic а surgery procedure as a result of growing public knowledge of cosmetic surgery. Liposuction is the process of employing a cannula and a strong vacuum to remove fat from deposits under the skin. Based on the amount of infiltration or wetting solution injected, four different liposuction procedures have been identified: dry, wet, superwet, and tumescent technique. The key distinction

between these procedures is how deeply they penetrate the tissues and how much blood is lost as a proportion of fluid sucked. The most popular method for liposuction is the tumescent method.

Tumescent technique

The phrase "tumescent method" was created by Klein. Tumescence literally means "to swell." In this procedure, very high quantities of diluted local anaesthetic are injected into the subcutaneous tissue combined with additions like adrenaline and sodium bicarbonate to expand the tissues and cause them to become hard, puffy, and turgid, or the ultimate goal of strong tissue turgor. As a result, a plane is created from which the suction of fat is made easier with minimal blood loss [1].

According to the amount of aspiration performed, there forms are two of liposuction: high volume (>4,000 ml aspirated) and low volume (4,000 ml aspirated). High-volume liposuction is seen as being substantially more prone to complications than the original form of tumescent liposuction since there is a possibility for major fluid changes related to the volume of tissue removed, with a risk of

Received: 09-November-2022, Manuscript No. ijocs-22-84477;

Editor assigned: 12-November-2022, PreQC No. ijocs-22-84477 (PQ);

Reviewed: 16-November-2022, QC No. ijocs-22-84477 (Q);

Revised: 19-November-2022, Manuscript No. ijocs-22-84477 (R);

Published: 23-November-2022, DOI:10.37532/1753-0431.2022.16(11). 272

⁺Author for correspondence: Jacob Taylor, Editorial Office, International Journal of Clinical Skills, London, United Kingdom, E-mail: ijclinicalskill@journalres.com

Short Communication

Taylor

developing pulmonary oedema. Therefore, in and offer intraoperative analgesia. As a result, high-volume liposuction procedures, intravenous fluids should be utilised with the utmost caution [2].

Local anaesthetics

The most widely used local anaesthetic in tumescent solutions is lidocaine, with a maximum dose of 7 mg/kg when combined with adrenaline. The maximum dose of lignocaine that can be administered in a tumescent solution may be much higher thanks to the development of tumescent anaesthesia [3-5]. For most patients, a maximum dose of 55 mg/kg of lignocaine is advised; those having liposuction should have a dose between 35 mg/kg and 55 mg/kg. The amount of lignocaine used depends on how vascular the area is that needs to be liposuctioned. It is possible to use 1,500 mg/L of normal saline in the breast and belly, which are more vascular or sensitive areas.

In less sensitive regions, such the thighs, the concentration of normal saline is lowered to 500 mg/L. Louis Habbema observed that a maximum lignocaine concentration of 500 mg/L of normal saline solution allowed infusion of a considerable volume of wetting solution without any risk of lignocaine poisoning in a series of 3,430 liposuction procedures.

Anaesthesia Technique

Under either local, regional, or general anaesthesia, liposuction can be carried out. There is no one anaesthetic method that has been shown to be better than the rest. The method of anaesthesia is determined by the location and extent of the liposuction as well as the patient's preferences. Depending on the body parts being operated on arms, thighs, abdomen, or buttocks and the amount of liposuction being done, the anaesthetic approach will change. Since these treatments are frequently carried out in a day care setting, it is critical that patients quickly regain their psychomotor and cognitive abilities to allow for early discharge.

Monitored anaesthesia care

This is a practical method for liposuction of tiny volumes. Vasoconstrictor and local anaesthetic infiltration, respectively, decrease bleeding it enables liposuction to be carried out under minimal sedation, resulting in a speedy recovery, earlier discharge, and lower patient costs. But if infiltration is uneven, certain places won't have enough analgesia, necessitating further sedation.

General anaesthesia

This is suggested for liposuction of high volumes or if the patient wants it. Due to its pharmacokinetic profile and innate antiemetic characteristic, which guarantees a speedy patient recovery, propofol is the preferred induction agent. A supraglottic device, proseal LMA (PLMA), or I-gel can be used to preserve airway. A nondepolarizing medication, such as atracurium, rocuronium, or vecuronium, can relax muscles. Short-acting opioids like fentanyl and remifentanil, as well as the injection of lignocaine into subcutaneous tissue, offer analgesia.

Postoperative Care

For large-volume liposuction, postoperative monitoring should continue.

Pain relief

In tumescent liposuction, pre-emptive analgesia is delivered by the local anaesthetic, so little intraoperative and postoperative analgesia is necessary. The amount of time that lignocaine provides analgesia can be greatly increased by mixing it with epinephrine. NSAIDs, often known as non-steroidal anti-inflammatory drugs, may be prescribed to treat postoperative pain.

Conclusion

A common method for getting rid of extra body fat is liposuction. It has a significant hemodynamic impact, particularly during large-volume liposuction operations. A full understanding of the physiological changes and potential consequences linked to them is necessary for the anaesthesia used for these procedures. A successful outcome is guaranteed by meticulous monitoring and rigorous adherence to intraoperative fluid treatment standards.

Short Communication

References

- 1. Maltseva N, Borzova E, Fomina D, et al. Cold urticarial-What we know and what we do not know Allergy. *Eur Acad Allergy Clin Immunol 76*, 1077-1094 (2021).
- 2. Maurer M, Metz M, Bindslev Jensen

C, et al. Definition, aims, and implementation of GA(2) LEN urticaria centers of reference and excellence Allergy. Eur Acad Allergy Clin Immunol 71, 1210-1218 (2016).

- 3. Ramos-Casals M, Stone JH, Cid MC, et al. The cryoglobulinaemias. *Lancet 379*, 348-360 (2012).
- 4. Bracken SJ, Abraham S, MacLeod AC. Autoimmune theories of chronic spontaneous urticarial. *Front Immunol 10*, (2019).
- 5. Koeppel MC, Bertrand S, Abitan R, et al. Urticaria caused by cold. *Ann Dermatol Venereol* 123, 627-632 (1996).